

Phonons and Solitons in 1D Mott Insulators

Chun-Min Chang¹, A. H. Castro Neto^{2,1}, and A. R. Bishop³

¹ *Department of Physics, University of California, Riverside, CA 92521*

² *Department of Physics, Boston University, Boston, MA 02215*

³ *Theoretical Division and Center for Nonlinear Studies, Los Alamos National Laboratory, Los Alamos, New Mexico 87545*

(February 1, 2008)

We study the problem of one-dimensional (1D) Luttinger liquids in the insulating Mott-Hubbard phase in the presence of acoustic phonons in the semiclassical limit. We show that solitonic excitations can propagate freely in the system and the energy required to produce a soliton is reduced by a polaronic effect. We find a critical value of the electron-phonon coupling constant for which this energy vanishes.

PACS numbers:71.10-Pm,71.38.-k

It is well-known that phonons can have a major impact in the phase diagram of 1D interacting electron systems, driving these systems from charge density waves (CDW) to superconductors due to retardation effects [1]. In fact, the problem of phonons in 1D has been studied to a great level of detail in the phase where the linear electronic charge density is incommensurate with the lattice leading to renormalizations of the various Luttinger liquid parameters and therefore to no dramatic effect in the charge propagation [2]. In this note we study the problem of the effect of acoustic phonons in the *commensurate* phase of Luttinger liquids when the 1D system has an instability to the opening of a gap (Mott gap) in the charge spectrum and therefore to insulating behavior. In the absence of phonons it can be shown that the lowest energy excitation is associated with the creation of a soliton (or anti-soliton) which can propagate freely in the system. This excitation, however, requires a finite amount of energy Δ_0 which is directly related to the Mott gap. As it is well-known the origin of the soliton is associated with the Umklapp scattering of electrons in the presence of the lattice [3]. In a bosonized description of this process one can show that the bosonic charge fields (associated with electronic charge excitations) are described by a sine-Gordon theory which has as semiclassical solutions traveling kinks and anti-kinks. Much less is known about effects of phonons on the soliton propagation in the insulating phase. We show that while the electron-phonon coupling allows for coherent soliton propagation in the Mott phase it can also lead to strong non-perturbative renormalization effects. We show here by a non-perturbative semiclassical calculation that the soliton excitation energy is reduced in the presence of acoustic phonons and that there is a critical value of the electron-phonon coupling for which the Δ , the gap for the soliton creation in the presence of phonons, vanishes. Physically, the electron-electron coupling is suppressed by a polaronic effect.

The basic electronic Hamiltonian we consider can be written in a very general form:

$$H_{el} = -t \sum_{i,\sigma,\alpha} c_{i,\sigma}^+ c_{i+1,\sigma} + \sum_{i,\sigma,\sigma'} V_{\sigma,\sigma'}(i,j) n_{i,\sigma} n_{j,\sigma'}, \quad (1)$$

where $c_{i,\sigma}$ ($c_{i,\sigma}^+$) is the annihilation (creation) electron operator at lattice site i , with spin projection $\sigma = \uparrow, \downarrow$ and $n_{i,\sigma} = c_{i,\sigma}^+ c_{i,\sigma}$ is the electron number operator. Here t is the kinetic energy and $V_{\sigma,\sigma'}(i,j)$ is the interaction energy between electrons on different sites. In what follows we assume short range interactions only. If, for instance, we keep only the on-site Coulomb repulsion $V_{\uparrow,\downarrow}(i,i) = U$ and the nearest neighbor Coulomb term $V_{\sigma,\sigma'}(i,i+1) = V$ this Hamiltonian describes the extended Hubbard Model. The many-body techniques we will apply here are actually valid for a very generic class of Hamiltonians [4]. As described elsewhere, [5] the fermionic degrees of freedom can be written in terms of bosonic charge, $\Phi_\rho(x)$, and spin, $\Phi_\sigma(x)$, fields. The bosonic version of the fermionic Hamiltonian (1) reads:

$$H_{el} = H_\rho + H_\sigma + \frac{2g_1}{(2\pi a)^2} \int dx \cos \left[\sqrt{8} \Phi_\sigma(x) \right] + \frac{2g_3}{(2\pi a)^2} \int dx \cos \left[\sqrt{8} \Phi_\rho(x) + 4k_F x \right], \quad (2)$$

where

$$H_v = \frac{1}{2} \int dx \left[(u_v K_v) \pi^2 \Pi_v^2(x) + \frac{u_v}{K_v} (\partial_x \Phi_v(x))^2 \right], \quad (3)$$

where $v = \rho, c$ refers to the free bosonic Hamiltonian (Π_v is the momentum field operator, that is, $[\Pi_v(x), \Phi_u(y)] = i\delta(x-y)\delta_{u,v}$ for charge and spin, respectively, with Luttinger liquid parameters K_v , velocities u_v , and coupling constants g_1 and g_3 that are directly related to the backscattering and Umklapp processes, respectively. Here $k_F = \pi n/2$ is the Fermi momentum where n is the linear density of electrons. All these parameters can be written in terms of the bare parameters of the original Hamiltonian (1). In this representation the electronic charge density can be written as:

$$\rho(x) = -\frac{\sqrt{2}}{\pi} \partial_x \Phi_\rho(x). \quad (4)$$

Observe that in (3) the charge and spin degrees of freedom do not interact with each other. This is a generic property of 1D quantum fluids (spin-charge separation). Many important results can be understood from the perturbative renormalization group (RG) studies of (2). In the presence of attractive electron-electron interactions the backscattering term, g_1 , is a relevant perturbation which leads to the opening of a gap in the spin spectrum which is associated with the dominance of superconducting fluctuations. In this case the charge excitations are gapless. In the presence of repulsive interactions backscattering is irrelevant and the Umklapp term, g_3 , is the most important one. When the electronic density, n , is commensurate with the lattice (that is, $n = 1/a$, where a is the lattice spacing) the $4k_F x$ term in (2) can be dropped and the Umklapp term becomes a relevant perturbation which leads to the opening of a gap in the charge spectrum. In this case the spin spectrum remains gapless. When n is incommensurate with the lattice both g_1 and g_3 are irrelevant and spin and charge both have gapless spectra.

Let us now consider the effect of an acoustic phonon field, $\phi(x)$, described by the Hamiltonian:

$$H_{ph} = \frac{1}{2} \int dx \left[P^2(x) / \rho_s + c_s^2 \rho_s (\partial_x \phi(x))^2 \right], \quad (5)$$

where $P(x)$ is the phonon momentum operator canonically conjugated to $\phi(x)$, ($[P(x), \phi(y)] = i\delta(x - y)$), $\rho_s = M/a$ is the lattice mass density and c_s is the speed of sound. As is well-known, [6] acoustic phonons couple to electrons via a deformation potential term which can be written as:

$$\begin{aligned} H_{el-ph} &= -\gamma \frac{\pi}{\sqrt{2}} \int dx \rho(x) \partial_x \phi(x) \\ &= \gamma \int dx \partial_x \Phi_\rho(x) \partial_x \phi(x), \end{aligned} \quad (6)$$

where we have used (4). Here γ is the electron-phonon coupling constant. Naively a perturbative RG analysis of (6) would indicate that the coupling is irrelevant since it has two derivatives. In fact, as we are going to show the electron-phonon coupling does not affect the soliton propagation (and therefore it is irrelevant from this perspective) but we are also going to show that the soliton energies are strongly renormalized by the polaronic effect generated by this operator and can even vanish in the strong coupling regime. Thus, the irrelevancy of the operator in the weak coupling regime does not imply that this operator is not important in the strong coupling limit.

Notice that the spin part of the Hamiltonian (3) does not couple to the phonons we consider and can be dropped. We focus now entirely on the charge degrees of freedom. When a charge is injected into the the Mott insulator it creates a soliton excitation which can be described in terms of the equation of motion for the bosonic

fields. Indeed, consider the equations of motion which are obtained from the Hamiltonian in the Heisenberg representation. It is very easy to show that the fields obey the following equations:

$$\begin{aligned} \frac{1}{u_\rho \pi K_\rho} \frac{\partial^2 \Phi_\rho}{\partial t^2} &= \frac{u_\rho}{\pi K_\rho} \frac{\partial^2 \Phi_\rho}{\partial x^2} + \frac{\sqrt{2} g_3}{(\pi a)^2} \sin(\sqrt{8} \Phi_\rho) + \gamma \frac{\partial^2 \phi}{\partial x^2} \\ \rho_s \frac{\partial^2 \phi}{\partial t^2} &= c_s^2 \rho_s \frac{\partial^2 \phi}{\partial x^2} + \gamma \frac{\partial^2 \Phi_\rho}{\partial x^2}. \end{aligned} \quad (7)$$

We now take advantage of the Lorentz invariance of these equations and make a change of variables $\lambda = x \pm vt$ where v is the soliton velocity, in order to write:

$$\begin{aligned} \left(\frac{v^2}{u_\rho \pi K_\rho} - \frac{u_\rho}{\pi K_\rho} \right) \frac{\partial^2 \Phi_\rho}{\partial \lambda^2} &= \frac{\sqrt{2} g_3}{(\pi a)^2} \sin(\sqrt{8} \Phi_\rho) + \gamma \frac{\partial^2 \phi}{\partial \lambda^2} \\ \rho_s (v^2 - c_s^2) \frac{\partial^2 \phi}{\partial \lambda^2} &= \gamma \frac{\partial^2 \Phi_\rho}{\partial \lambda^2}. \end{aligned} \quad (8)$$

Combining both equations and replacing Φ_ρ with $\Phi = \sqrt{8} \Phi_\rho$, the equation of motion for the charge field is given by:

$$\mu \frac{\partial^2 \Phi}{\partial \lambda^2} + \sin \Phi = 0, \quad (9)$$

which is a sine-Gordon equation with

$$\mu = \frac{(\pi a)^2}{4g_3} \left[\frac{\gamma^2}{\rho_s (c_s^2 - v^2)} - \frac{(u_\rho^2 - v^2)}{u_\rho \pi K_\rho} \right]. \quad (10)$$

As is well-known this equation has single soliton solutions:

$$\begin{aligned} \Phi_\rho(x, t) &= \pm \sqrt{2} \tan^{-1} \left(e^{\frac{1}{\sqrt{|\mu|}}(x \pm vt)} \right), \quad \text{for } \mu > 0 \\ &= \pm \sqrt{2} \tan^{-1} \left(e^{\frac{1}{\sqrt{|\mu|}}(x \pm vt)} \right) + \pi, \quad \text{for } \mu < 0. \end{aligned} \quad (11)$$

Moreover, from (8), we find that the lattice deformation also propagates coherently and has a profile given by

$$\phi(x, t) = \pm \frac{\sqrt{2} \gamma}{\rho_s (v^2 - c_s^2)} \tan^{-1} \left(e^{\frac{1}{\sqrt{|\mu|}}(x \pm vt)} \right). \quad (12)$$

These equations show that coherent charge propagation is possible in the form of a soliton (or kink dressed with a correlated lattice deformation). The minimum energy required to create such an excitation can be obtained by substituting the above solutions for the boson and phonon field directly into the Hamiltonian. After straightforward algebra we find that the energy required for soliton creation is given by:

$$\Delta = \frac{2}{\pi a} \sqrt{\left| g_3 \left(\frac{u_\rho}{\pi K_\rho} - \frac{\gamma^2}{\rho_s c_s^2} \right) \right|},$$

which is reduced from its bare value $\Delta_0 = \frac{2}{\pi a} \sqrt{g_3 u_\rho / (\pi K_\rho)}$ by an amount which depends on the electron-phonon coupling constant. So far this calculation is semiclassical in nature since it does not consider quantum corrections which will be discussed elsewhere [7]. This result predicts, however, that Δ vanishes when γ attains a critical value

$$\gamma_c = c_s \sqrt{\frac{u_\rho \rho_s}{\pi K_\rho}}. \quad (13)$$

We thus predict that in systems with strong electron-phonon coupling the energy required to create a soliton vanishes due to the polaronic effect described here. Note that in the absence of the Umklapp term arising from the discrete lattice the effects described here are not possible.

This effect might explain the recently observed long range charge transport in DNA (which we claim is a Mott insulator [8]) where injected charge carriers travel long distances along the DNA double helix [9].

We thank illuminating discussions with J. Barton, W. Beyermann, D. Cox, G. Gruner, and S. Kivelson. We especially thank M. Pollak for stimulating our interest in this problem. We acknowledge partial support provided by the Collaborative University of California - Los Alamos (CULAR) research grant under the auspices of the US Department of Energy.

-
- [1] G. T. Zimanyi, S. A. Kivelson and A. Luther, Phys. Rev. Lett. **60**, 2089 (1988); S. Kivelson, and G. T. Zimanyi, Mol. Cryst. Liq. Cryst. **160**, 457 (1988).
 - [2] T. Martin and D. Loss, Int. J. Mod. Phys. **9**, 495 (1995); O. Heinonen and S. Eggert, Phys. Rev. Lett **77**, 358 (1996).
 - [3] T. Giamarchi, Physica B **230**, 975 (1997).
 - [4] V. J. Emery, in *Highly Conducting One-Dimensional Solids*, Edited by J. T. Devreese, R. P. Evrard and V. E. van Doren, Plenum Press (1979); J. Sólyom, "The Fermi gas model of one-dimensional conductors", Adv.Phys. **28**, 201, (1979).
 - [5] For a review, see, J. Voit, Rep. Prog. Phys. **58**, 977 (1995).
 - [6] G. Mahan in *Many-Particle Physics* (Plenum Press, New York 1990).
 - [7] Chun-Min Chang, A. H. Castro Neto, and A. R. Bishop, in preparation.
 - [8] Chun-Min Chang, A. H. Castro Neto, and A. R. Bishop, cond-mat/0008166.
 - [9] D. B. Hall, R. E. Holmlin and J. K. Barton, Nature **382**, 731 (1996); S. O. Kelley and J. K. Barton, Science **283**, 375 (1999).